

2013

A Simple Chelex Protocol for DNA Extraction From Anopheles Spp

M. Musapa

T. Kumwenda

M. Mkulama

S. Chishimba

D. E. Norris

See next page for additional authors

Follow this and additional works at: https://mosaic.messiah.edu/bio_ed



Part of the [Biology Commons](#)

Permanent URL: https://mosaic.messiah.edu/bio_ed/105

Recommended Citation

Musapa, M.; Kumwenda, T.; Mkulama, M.; Chishimba, S.; Norris, D. E.; Thuma, Philip; and Mharakurwa, S., "A Simple Chelex Protocol for DNA Extraction From Anopheles Spp" (2013). *Biology Educator Scholarship*. 105.

https://mosaic.messiah.edu/bio_ed/105

Sharpening Intellect | Deepening Christian Faith | Inspiring Action

Messiah University is a Christian university of the liberal and applied arts and sciences. Our mission is to educate men and women toward maturity of intellect, character and Christian faith in preparation for lives of service, leadership and reconciliation in church and society.

Authors

M. Musapa, T. Kumwenda, M. Mkulama, S. Chishimba, D. E. Norris, Philip Thuma, and S. Mharakurwa

Video Article

A Simple Chelex Protocol for DNA Extraction from *Anopheles* spp.

Mulenga Musapa¹, Taida Kumwenda¹, Mtawa Mkulama¹, Sandra Chishimba¹, Douglas E. Norris², Philip E. Thuma¹, Sungano Mharakurwa¹

¹Malaria Institute at Macha

²Department of Molecular Microbiology & Immunology, Johns Hopkins Bloomberg School of Public Health

Correspondence to: Sungano Mharakurwa at sungano.mharakurwa@miam.org.zm

URL: <http://www.jove.com/video/3281>

DOI: [doi:10.3791/3281](https://doi.org/10.3791/3281)

Keywords: Infection, Issue 71, Immunology, Infectious Diseases, Genetics, Molecular Biology, Microbiology, Parasitology, Entomology, Malaria, *Plasmodium falciparum*, vector, *Anopheles*, Diptera, mosquitoes, Chelex, DNA, extraction, PCR, dissection, insect, vector, pathogen

Date Published: 1/9/2013

Citation: Musapa, M., Kumwenda, T., Mkulama, M., Chishimba, S., Norris, D.E., Thuma, P.E., Mharakurwa, S. A Simple Chelex Protocol for DNA Extraction from *Anopheles* spp.. *J. Vis. Exp.* (71), e3281, doi:10.3791/3281 (2013).

Abstract

Endemic countries are increasingly adopting molecular tools for efficient typing, identification and surveillance against malaria parasites and vector mosquitoes, as an integral part of their control programs^{1,2,3,4,5}. For sustainable establishment of these accurate approaches in operations research to strengthen malaria control and elimination efforts, simple and affordable methods, with parsimonious reagent and equipment requirements are essential^{6,7,8}. Here we present a simple Chelex-based technique for extracting malaria parasite and vector DNA from field collected mosquito specimens.

We morphologically identified 72 *Anopheles gambiae* *sl.* from 156 mosquitoes captured by pyrethrum spray catches in sleeping rooms of households within a 2,000 km² vicinity of the Malaria Institute at Macha. After dissection to separate the head and thorax from the abdomen for all 72 *Anopheles gambiae* *sl.* mosquitoes, the two sections were individually placed in 1.5 ml microcentrifuge tubes and submerged in 20 μ l of deionized water. Using a sterile pipette tip, each mosquito section was separately homogenized to a uniform suspension in the deionized water. Of the ensuing homogenate from each mosquito section, 10 μ l was retained while the other 10 μ l was transferred to a separate autoclaved 1.5 ml tube. The separate aliquots were subjected to DNA extraction by either the simplified Chelex or the standard salting out extraction protocol^{9,10}. The salting out protocol is so-called and widely used because it employs high salt concentrations in lieu of hazardous organic solvents (such as phenol and chloroform) for the protein precipitation step during DNA extraction⁹.

Extracts were used as templates for PCR amplification using primers targeting arthropod mitochondrial nicotinamide adenine dinucleotide dehydrogenase (NADH) subunit 4 gene (ND4) to check DNA quality¹¹, a PCR for identification of *Anopheles gambiae* sibling species¹⁰ and a nested PCR for typing of *Plasmodium falciparum* infection¹². Comparison using DNA quality (ND4) PCR showed 93% sensitivity and 82% specificity for the Chelex approach relative to the established salting out protocol. Corresponding values of sensitivity and specificity were 100% and 78%, respectively, using sibling species identification PCR and 92% and 80%, respectively for *P. falciparum* detection PCR. There were no significant differences in proportion of samples giving amplicon signal with the Chelex or the regular salting out protocol across all three PCR applications. The Chelex approach required three simple reagents and 37 min to complete, while the salting out protocol entailed 10 different reagents and 2 hr and 47 min' processing time, including an overnight step. Our results show that the Chelex method is comparable to the existing salting out extraction and can be substituted as a simple and sustainable approach in resource-limited settings where a constant reagent supply chain is often difficult to maintain.

Video Link

The video component of this article can be found at <http://www.jove.com/video/3281/>

Protocol

1. Preparatory Dissection of Mosquito Specimens

1. Place mosquito on a small cutting of parafilm and mount on dissecting microscope.
2. Cover in drop of deionized water to soften tissue.
3. Incise the mosquito carcass precisely at the joint between the thorax and abdomen, thus nicking off the head and thorax from the abdominal section.
4. Transfer each dissected mosquito section into a clean autoclaved 1.5 ml microcentrifuge tube pre-labeled with mosquito ID number details and appropriately marked to denote body section (e.g. "m" for mid-gut section; "h" for head and thorax section).
5. Discard parafilm and carefully wipe microscope stage with 70% alcohol-moistened Kimwipe before processing the next mosquito.

2. DNA Extraction from Mosquito Specimens

1. Pipette 20 μ l of deionized water into sample tube and use pipette tip to grind the submerged mosquito section into a uniform suspension.
2. Add 100 μ l of autoclaved 1X PBS/1% saponin solution to sample homogenate and mix by gentle momentary vortexing.
3. Incubate at room temperature for 20 min.
4. Centrifuge at 20,000 x g for 2 min and discard supernatant.
5. Resuspend pellet in 100 μ l 1X PBS.
6. Centrifuge again at 20,000 x g for 2 min and discard supernatant.
7. By gentle vortexing (5 sec), resuspend pellet in 75 μ l sterile deionized water and 25 μ l of 20% w/v Chelex-100 resin suspension in deionized water.
8. Pierce hole in lid of sample tube using sterile 23G hypodermic needle flamed in Bunsen burner.
9. Boil sample suspension in water bath on floating rack (or in heating block) for 10 min.
10. Centrifuge at 20,000 x g, for 1 min and transfer ensuing DNA solution into prelabeled storage vial for use as template in PCR applications.

3. Plasmodium falciparum Genotyping and Anopheles spp. Molecular Identification

1. Add 2.5 μ l of the Chelex DNA extract in 25 μ l PCR reactions to check DNA quality¹¹, identify *Anopheles* species¹⁰ and to genotype *P. falciparum*¹² mid-gut and salivary gland infections.
2. To maximize amplicon yield, especially for *P. falciparum* detection, include 1.5X bovine serum albumin (BSA) in the PCR assay. The following reaction composition is recommended: (2.5 μ l template, 0.25 μ M primers, 1.5 μ M magnesium chloride, 200 μ M dNTPs, 1X PCR Buffer, 1.5X BSA and 1.0U Taq DNA polymerase, in 25 μ l volumes).
3. To check for DNA quality in extract, amplify region of the arthropod mitochondrial nicotinamide adenine dinucleotide dehydrogenase (NADH) subunit 4 (ND4) gene (primers ND4FW [5'-GTD YAT TTA TGA TTR CCT AA-3'] and ND4RV [5'-CTT CGD CTT CCW ADW CGT TC-3']; expect product 400bp)^{11,13}.
4. To differentiate *Anopheles gambiae* *sl.* sibling species (*An. gambiae* *ss* and *An. arabiensis* only), amplify region flanking SNPs in the intergenic spacer (IGS) region, (primers UN [5'-GTGTGCCCTTCCTCGATGT-3'], GA [5'-CTGGTTTGGTCGGCACGTTT-3'], AR [5'-AAGTTCCTTCTCCATCCTA-3']; expect product 390bp *An gambiae* *ss*, 315bp *An arabiensis*)^{10,11}.
5. For typing *P. falciparum* antifolate drug resistance polymorphisms, perform nested PCR to amplify region flanking amino acid codons 108, 51, 59, 16 and 164 in the parasite dihydrofolate reductase (DHFR) gene:
Primary round primers: M1 [5'-TTTATGATGGAACAAGTCTGC-3'] and M5 [5'-AGTATATACATCGCTAACAGA-3']
Secondary round primers: M3 [5'-TTTATGATGGAACAAGTCTGCGACGTT-3'] with F/ [5'-AAATTCTTGATAAACAACGGAAACCTTTTA-3']
 Or
 F [5'-GAAATGTAATTCCTAGATATGGAATATT-3'] with M4 [5'-TTAATTTCCCAAGTAAACTATTAGAGCTTC-3'].
 Also amplify region containing amino acid codons 436, 437, 540, 581 and 613 in the *P. falciparum* dihydropteroate synthetase (DHPS) gene:
Primary round primers R2 [5'- AACCTAAACGTGCTGTTCAA-3'] with R/ [5'- AATTGTGTGATTTGTCCACAA3'];
Secondary round primers
 J: [5'-TGCTAGTGTATAGATATAGGTGGAGAAAGC-3'] with K/ [5'-CTATAACGAGGTATTGCATTTATTGCAAGAA-3']
 Or
 K: [5'-TGCTAGTGTATAGATATAGGATGAGCATC-3'] with K/,
 Or
 L [5'- ATAGGATACTATTGATATTGATATTGGACCAGGATTCG-3'] with L/ [5'-5TATTACAACATTTTGATCATTGCGCAACCGG-3']¹².
6. Subject 4 μ l amplicon to allele-specific restriction enzyme digestions in 30 μ l reactions following the manufacturer's instructions, to detect antifolate drug resistance-associated polymorphisms¹².
7. Subject 5 μ l of PCR amplicon (or 15 μ l of restriction digest) per lane of ethidium bromide-stained 2% agarose gel to electrophoresis (100 - 120V) and visualize bands under UV transillumination.

Representative Results

Examples of results from PCR assays for mosquito DNA extract quality (**Figure 1**), *Anopheles arabiensis* molecular identification (**Figure 2**) and *P. falciparum* detection (**Figure 3**) show that the simplified Chelex method yields similar results to the standard salting out protocol¹⁰, despite much fewer steps (**Table 1**). With comparable DNA quality in the respective extracts it is not surprising that sample positivity rates with respect to *An. gambiae* sibling species as well as parasite infection rates were not statistically different based on McNemar's chi-square test (**Figure 3**).

Sensitivity (%) was calculated as TP/(TP+FN)*100, where TP denotes true positives and FN denotes false negatives. Specificity (%) was determined as TN/(TN+FP)*100, where TN denotes true negatives and FP denotes false positives. The DNA quality (ND4) PCR showed 93% sensitivity and 82% specificity for the Chelex approach compared to the established salting out protocol as gold standard. Corresponding values of sensitivity and specificity were 100% and 78%, respectively, using sibling species identification PCR and 92% and 80%, respectively for *P. falciparum* detection PCR.

The addition of BSA to reaction mixtures resulted in a general increase of PCR positives (**Figure 3**) due to relief of PCR inhibitors¹⁴, for both the simplified Chelex and regular salting out protocol. However, this increase was not statistically significant except for DNA quality (ND4 PCR) on Chelex extracts ($p = 0.039$). Allele-specific restriction enzyme digestion on *P. falciparum* DHFR (or other target gene) amplicon enables the genotyping of mid-gut and salivary gland malaria infections for drug resistance alleles (**Figure 4; salivary gland data not shown**).

Simple Chelex Procedure		Standard Salting out Procedure	
Steps	Reagents	Steps	Reagents

<ol style="list-style-type: none"> 1. Add specimen to 1.5 ml microfuge tube and homogenize in 100 μl of 1X PBS/1% Saponin solution (2 min). 2. Leave at room temperature for 20 min. 3. Spin at 20,000 x g for 2 min and discard supernatant. 4. Add 100 μl of 1X PBS and mix gently by momentarily vortexing (3 sec). 5. Spin at 20,000 x g for 2 min and discard supernatant. 6. Add 25 μl of 20% w/v Chelex in deionized water and 75 μl sterile deionized water. 7. Pierce hole in lid of sample tube using hot, sterile 23G hypodermic needle and boil tube contents for 10 min. 8. Spin microfuge tube at 20,000 x g for 1 min. 9. Transfer supernatant into sterile prelabeled vial and store DNA solution at -20 $^{\circ}$C until use (2.5 μl in 25 μl PCR reaction). 	<p>PBS Saponin Chelex-100 beads</p>	<ol style="list-style-type: none"> 1. Add specimen to 1.5 ml microfuge tube and homogenize in 100 μl Bender buffer* with 1% DEPC (2 min). 2. Incubate at 65 $^{\circ}$C for 1 hr. 3. Add 15 μl cold 8 M potassium acetate. Mix gently and incubate on ice for 45 min. 4. Spin sample in microfuge tubes at 20,000 x g for 10 min and transfer supernatant to a new 1.5 ml tube. 5. Add 250 μl of 100% ethanol and mix well by inverting the tube. 6. Incubate sample at RT for 5 min. 7. Spin samples at 20,000 x g for 15 min. 8. Aspirate and discard supernatant, leaving the pellet to dry completely in the tube (30 min). 9. Resuspend pellet in 20 μl of 0.1X SSC + RNase (10 μg/ml) overnight at 4 $^{\circ}$C. 10. Add 80 μl DEPC water and store at -20 $^{\circ}$C until use. 	<p>Diethylpyrocarbonate (DEPC) *Bender Buffer: 0.1 M NaCl 0.2 M Sucrose 0.1 M Tris-HCL 0.05 M EDTA, pH 9.1 0.5% SDS in DEPC water 8 M Potassium acetate Ethanol 0.1X Saline sodium citrate (SSC) buffer RNase (10 μg/ml)</p>
<p>TOTAL TIME: 37 min</p>	<p>TOTAL TIME: 2 hr 47 min, plus overnight</p>		

Table 1. Step by step comparison of the reagents and time requirements for simple Chelex protocol and standard salting out method for DNA extraction from mosquito specimens.




Figure 1. ND4 mitochondrial PCR comparison of DNA quality from simplified Chelex "C" and standard salting out "M" extracts on *Anopheles arabiensis* field samples. NC, negative control; M1 and C1, M2 and C2, M3 and C3, and M4 and C4 are paired salting out and Chelex extracts from same *An. arabiensis* mosquito specimens. L, 100 bp DNA ladder.




Figure 2. Molecular identification PCR for *Anopheles arabiensis*. C1 and M1, C2 and M2, C3 and M3, C4 and M4 denote respective amplicon from salting out "M" and simplified Chelex "C" DNA extracts for same mosquito specimens; AR, *Anopheles arabiensis* positive control; L, 100 bp DNA ladder.




Figure 3. Detection of positives on Chelex and salting out DNA extracts for mosquito field samples, with PCR assays for molecular identification of *Anopheles arabiensis* (AR)¹⁰, arthropod NADH dehydrogenase gene 4 (ND4) DNA quality test¹¹, and antifolate drug resistance *P. falciparum* DHFR genotyping¹² (F-M4). Results shown for assays run with or without BSA in the reaction mix. McNemar's chi-square test was used to determine if the differences between percent of positive PCRs from DNA extracted from the simplified Chelex and the standard salting out protocols were statistically significant.




Figure 4. Application of simplified Chelex protocol in genotyping *P. falciparum* infections in mosquitoes (mid-gut data shown) for drug resistance alleles. BstN I digestion on amplicon flanking codon 108 of the *P. falciparum* DHFR gene¹² shows cycloguanil-resistant S108T mutants in mosquito samples (M1-M5; mid-gut data shown). U, undigested 522 bp amplicon; FCR3, laboratory standard *P. falciparum* positive control clone carrying S108T; K1, *P. falciparum* laboratory standard clone negative control carrying cycloguanil-sensitive S108N; L, 100 bp DNA ladder.

Discussion

The simplified Chelex method presented here enables extraction of quality *Anopheles spp* and *P. falciparum* DNA from mosquito specimens amenable to diverse PCR applications. This technique can be employed for molecular identification of malaria vector mosquitoes and surveillance of drug-resistant *P. falciparum* alleles in mosquitoes for national malaria control programs. The advantages of the simplified Chelex technique include simplicity, fewer reagents and hence cost, safety and shorter processing time (37 min) than standard protocols such as the salting out method¹⁰ (2 hr 47 min and an overnight step, **Table 1**). The aforementioned advantages and minimal reagent requirements (3 reagents) compared to the current standard protocol (10 reagents, **Table 1**)^{11,15}, make the simplified Chelex protocol particularly friendly to endemic country laboratories where a constant reagent supply chain is often difficult to maintain. A limitation of the method is that like the standard protocol, it was also subject to PCR inhibitors known to occur in the mosquito integument¹⁶. However, this is readily relieved by inclusion of BSA in the assays. BSA has also been successfully employed as an amplification enhancer against inhibitors in other PCR applications^{17,18}.

Disclosures

No conflicts of interest declared.

Acknowledgements

The authors are indebted to the chiefs, headmen and communities of Macha for their unwavering cooperating during mosquito field sample collections. Dr Doug Norris and Rebekah Kent offered invaluable expertise on the salting out protocol. This work was funded by the Johns Hopkins Malaria Research Institute pilot grant system. Mosquito and parasite DNA laboratory standards were kindly donated by MR4, American Type & Culture Collection.

References

1. Bonnet, M., *et al.* Efficacy of antimalarial treatment in Guinea: *in vivo* study of two artemisinin combination therapies in Dabola and molecular markers of resistance to sulphadoxine-pyrimethamine in N'Zerekore. *Malar. J.* **6**, 54, [pii] 1475-2875-6-54 doi:10.1186/1475-2875-6-54 (2007).

2. Nsimba, B., *et al.* Sulphadoxine/pyrimethamine versus amodiaquine for treating uncomplicated childhood malaria in Gabon: a randomized trial to guide national policy. *Malar. J.* **7**, 31, [pii] 1475-2875-7-31 doi:10.1186/1475-2875-7-31 (2008).
3. Oyewole, I.O., *et al.* Behaviour and population dynamics of the major anopheline vectors in a malaria endemic area in southern Nigeria. *J. Vector Borne Dis.* **44**, 56-64 (2007).
4. Harris, I., *et al.* A large proportion of asymptomatic Plasmodium infections with low and sub-microscopic parasite densities in the low transmission setting of Temotu Province, Solomon Islands: challenges for malaria diagnostics in an elimination setting. *Malar. J.* **9**, 254, [pii] 1475-2875-9-254 doi:10.1186/1475-2875-9-254 (2010).
5. Ouedraogo, A.L., *et al.* Substantial contribution of submicroscopical *Plasmodium falciparum* gametocyte carriage to the infectious reservoir in an area of seasonal transmission. *PLoS ONE.* **4**, e8410, doi:10.1371/journal.pone.0008410 (2009).
6. Birtz, D., de Souza, M., & Nkengasong, J.N. Laboratory challenges in the scaling up of HIV, TB, and malaria programs: The interaction of health and laboratory systems, clinical research, and service delivery. *Am. J. Clin. Pathol.* **131**, 849-851, doi:131/6/849 [pii] 10.1309/AJCPGH89QDSWFONS (2009).
7. Ishengoma, D.R., *et al.* Health laboratories in the Tanga region of Tanzania: the quality of diagnostic services for malaria and other communicable diseases. *Ann. Trop. Med. Parasitol.* **103**, 441-453, doi:10.1179/136485909X451726 (2009).
8. Alonso, P.L., *et al.* A research agenda for malaria eradication: vector control. *PLoS Med.* **8**, e1000401, doi:10.1371/journal.pmed.1000401 (2011).
9. Grimberg, J., *et al.* A simple and efficient non-organic procedure for the isolation of genomic DNA from blood. *Nucleic Acids Res.* **17**, 8390 (1989).
10. Scott, J.A., Brogdon, W.G., & Collins, F.H. Identification of single specimens of the *Anopheles gambiae* complex by polymerase chain reaction. *American Journal of Tropical Medicine and Hygiene.* **49**, 520-529 (1993).
11. Kent, R.J. & Norris, D.E. Identification of mammalian blood meals in mosquitoes by a multiplexed polymerase chain reaction targeting cytochrome B. *Am. J. Trop. Med. Hyg.* **73**, 336-342, [pii] 73/2/336 (2005).
12. Duraisingh, M.T., Curtis, J., & Warhurst, D.C. *Plasmodium falciparum*: detection of polymorphisms in the dihydrofolate reductase and dihydropteroate synthetase genes by PCR and restriction digestion. *Exp. Parasitol.* **89**, 1-8 (1998).
13. Gorrochotegui-Escalante, N., Munoz, M.L., Fernandez-Salas, I., Beaty, B.J., & Black, W.C.T. Genetic isolation by distance among *Aedes aegypti* populations along the northeastern coast of Mexico. *Am. J. Trop. Med. Hyg.* **62**, 200-209 (2000).
14. Kreader, C.A. Relief of amplification inhibition in PCR with bovine serum albumin or T4 gene 32 protein. *Appl. Environ. Microbiol.* **62**, 1102-1106 (1996).
15. Norris, D.E., Shurtleff, A.C., Toure, Y.T., & Lanzaro, G.C. Microsatellite DNA polymorphism and heterozygosity among field and laboratory populations of *Anopheles gambiae* ss (Diptera: Culicidae). *J. Med. Entomol.* **38**, 336-340 (2001).
16. Schriefer, M.E., Sacci, J.B., Jr., Wirtz, R.A., & Azad, A.F. Detection of polymerase chain reaction-amplified malarial DNA in infected blood and individual mosquitoes. *Exp. Parasitol.* **73**, 311-316, [pii] 0014-4894(91)90102-3 (1991).
17. Al-Soud, W.A. & Radstrom, P. Purification and characterization of PCR-inhibitory components in blood cells. *J. Clin. Microbiol.* **39**, 485-493, doi:10.1128/JCM.39.2.485-493.2001 (2001).
18. Hyun, C., Filippich, L.J., & Hughes, I. The inhibitory effect of pentobarbitone on reverse transcription-PCR. *J. Biochem. Biophys. Methods.* **62**, 63-68, [pii] S0165-022X(04)00185-X doi:10.1016/j.jbbm.2004.09.005 (2005).